Molecular characterization of *Pseudomonas aeruginosa* isolates from various clinical specimens in Khartoum/Sudan: Antimicrobial resistance and virulence genes

Islam A. Babour¹,
Maha B. Mohamed²,
Asem A. Shehabi¹

- ¹ Department of Pathology-Microbiology and Forensic Medicine, The University of Jordan, School of Medicine, Amman, Jordan.
- ² Department of Microbiology, The University of Medical Sciences and Technology, Faculty of Medical Laboratory Sciences, Khartoum.

Abstract

Background: *Pseudomonas aeruginosa* is a pathogenic organism responsible for frequent wound and nosocomial infections worldwide. Its infections are difficult to control since the organism is known to rapidly develop antibiotic resistance and becomes multidrug-resistant (MDR) during treatment of patients.

Aim of the study: This study was intended to investigate the occurrence of certain important types of (ESBL) and (MBL) enzymes in association with important specific virulence factors associated with *P. aeruginosa* clinical isolates from Khartoum, Sudan.

Methods: This study investigated 70 *P. aeruginosa* isolates which were collected from patients admitted to four major hospitals in Khartoum (Fedail, Ribat, Ibn Sina and Soba hospitals). These isolates were recovered from 40 wound swabs (57.1%), 27 urine samples (38.6%), and 3 pleural fluid samples (4.3%) of patients. Higher numbers of isolates were recovered from males 42 (60%) than in females 28 (40%). All *P. aeruginosa* isolates were first confirmed by conventional biochemical and second using molecular PCR tests. PCR methods were also used for detecting the presence of the virulence genes *ToxA*, *AlgD*, *LasB*, *exoS*, *exoU*, *CTX*, *GES-1*, and genes of *VIM*, *IMP*, *KPC*, *CTX*, *VEB-1* and *SHV-1*.

Results: Antimicrobial susceptibility testing of *P. aeruginosa* isolates showed a high resistance to azetronam 49 (70%), followed by ceftazidime 32 (45.7%), 16 ciprofloxacin (22.9%), gentamicin 13 (18.6%),

Contact information:

Islam A. Babour.

✉ esooo99@gmail.com

piperacillin-tazobactam 11 (15.7%), amikacin 9 (12.9%), and imipenem 6 (8.6%) showed the least resistance. All isolates were positive for *algD* and *lasB* (100%), followed by *toxA* (90%), *exoS* (34.3%), *exoU* (24.3%), respectively. The rates of detected ESBL genes *blaTEM*, *blaCTX-m*, *blaSHV-1*, *GES-1*, were 3.3%, 6.6%, 10%, 3.3%, 10%, respectively, but all isolates were negative for *bla-KPC* and *bla-VIM* and *IMP*. The percentages of pigment production were 61.4% for *pyocyanin*, 37.1% for *pyoverdin* and 1.4% for *pyorubin*.

Conclusion: The study demonstrated high rates of antimicrobial resistance markers to most commonly used antibiotics in treatment of *P. aeruginosa* infections. The majority of the isolates from urine and wound samples carried at least three potential virulence factor genes of *algD*, *lasB* and *toxA* and without any significant relation to their antimicrobial resistance markers.

Keywords

Pseudomonas Aeruginosa; Antimicrobial Resistance; Virulence Factors; Khartoum; Sudan.

Received 11-02-2020; Accepted 28-02-2020

Introduction

Pseudomonas aeruginosa is an opportunistic pathogen. The organism is frequently causes nosocomial infection, wound sepsis, pneumonia and urinary tract infections (UTIs) in hospitalized patients, especially in intensive care units (ICUs) [1-5].

P. aeruginosa needs very limited nutritional requirements. It can survive in distilled water [6]. The organism survives at wide range of temperature (4-42 °C), and it is also tolerant to disinfectants and high concentrations of salts [7]. *P. aeruginosa* is capable of producing several different pigments, which are pyocyanin (blue), pyoverdin (yellow and fluorescent), pyorubin (red), and pyomelanin (brown) [8].

Multidrug-resistant *P. aeruginosa* is considered to be a serious problem in all types of infections because the organism has intrinsic resistance to several drug classes and it is able to acquire resistance

to most normally effective anti-Pseudomonal drugs within a short period [9].

There are many virulence factors that may contribute to pathogenic potential of *P. aeruginosa*, such as elastase, alkaline protease hemolysins (phospholipase and lecithinase) and pyocyanin diffusible pigment. In addition, most *P. aeruginosa* strains produce a variety of toxins especially exotoxin encoded by the *toxA* gene, which inhibits protein biosynthesis and is responsible for tissue necrosis. Exoenzyme S is encoded by the *exoS* gene, which is secreted by a type-III secretion system directly into the cytosol of host epithelial cells and subsequently causes their destruction [10]. Other important virulence factor found in *P. aeruginosa* include *algD*, which codes for GDP mannose dehydrogenase, which is transcriptionally activated in mucoid strains, and elastase B, a zinc metallopro-

tease encoded by the (*lasB*) gene, which attacks eukaryotic cells [11-12]. *P. aeruginosa* may also possess flagellae and pili, which helps in motility and adherence to host cells [13].

In recent years, a new variety of β -lactamases have been detected in *P. aeruginosa* clinical isolates. Broad spectrum β -lactam resistance attributed to β -lactamases is generally due to the metallo- β -lactamases (MBLs) such as IMP and VIM. The over expression of chromosomally encoded cephalosporinase, AmpC, is prevalent in *P. aeruginosa*. These MBLs hydrolyze anti-Pseudomonal cephalosporins and carbapenems effectively and their activity is not suppressed by the β -lactamase inhibitors that are currently available for treatment [14-15].

This study aimed to investigate *P. aeruginosa* isolates obtained from various clinical specimens of patients in Khartoum, Sudan, for their antimicrobial resistance profile, the occurrence of most important MBL genes and the common specific virulence factors using PCR molecular and culture tests.

Materials and Methods

This study included a total of 70 different clinical *P. aeruginosa* isolates collected from patients admitted to four hospitals in Khartoum, Sudan, during the period between September to November 2017. All patients were on treatment with various antibiotics.

The data about each patient was obtained and registered as the following: name, gender, age, and type of specimen, and all patients are agreed for any necessary medical process once they are admitted to each hospital.

Culture and isolation

All *P. aeruginosa* isolates were preserved in cryo tubes containing brain heart infusion with 15% glycerol samples and transported from Sudan to Jordan. All isolates were then subcultured on *Pseudomonas* Agar plates and incubated for 24 to 48 hours at 37

°C, and were examined for the presence of pigmentation and grape-like odor of *P. aeruginosa* at the research laboratories of the School of Medicine, The University of Jordan, Amman, Jordan.

Identification and storage

All growth of *P. aeruginosa* isolates was first identified using conventional biochemical tests including oxidase test, develop pigmentation and growth on 42 °C and produce no chemical reaction during culture in Kligler's iron agar tube were stored in cryo tubes at -70°C.

Antimicrobial susceptibility using disc diffusion method

Antimicrobial susceptibility using disc diffusion test was performed according to the guidelines of the Clinical Laboratory and Standards Institute (CLSI, 2016) [16].

Minimum inhibitory concentration (MIC) using E-test

This test was only performed on the *P. aeruginosa* isolates, which were MDR to three or more antibiotic classes. Four classes of antibiotics were tested (ceftazidime, ciprofloxacin, imipenem and amikacin). Results were interpreted according to the CLSI 2016 guidelines [16].

DNA extraction and PCR

The DNA extraction was performed according to the Promega kit (USA). Two PCR assays were performed to confirm the isolates; one is specific for the genus *Pseudomonas*, while the other is specific for *P. aeruginosa*. Two pairs of primers were used for each assay based on 16S ribosomal DNA (rDNA) sequence as shown in **Table 1**.

The bacterial plasmid of freshly cultured *P. aeruginosa* isolates was extracted according to manufacturer instructions using the Zyppy™ Plasmid Miniprep Kit (QIAGEN, Germantown, MD, USA) for detection of ESBLs, MBLs and virulence genes. All

Table 1. 16S rDNA primers sets for differentiation of *P. aeruginosa* from *Pseudomonas* species and their sequences and product size.

Target	Primer	Primer sequence (5' to 3')	Product size(bp)	Reference
<i>Pseudomonas species</i>	PA-GS-F PA-GS-R	GACGGGTGAGTAATGCCTA CACTGGTGTTCCTTCTATA	618	17
<i>P. aeruginosa</i>	PA-SS-F PA-SS-R	GGGGGATCTTCGGACCTCA TCCTTAGAGTGCCACCCG-	956	17

Table 2. Primer targets, sequences and their product size for five ESBLs and virulence genes.

Target	Primer	Primer sequence (5' to 3')	Product size (bp)	Reference
blaCTX-M	CTX-M(F) CTX-M(R)	CGCTTTGCGATGTGCAG ACCGCGATATCGTTGGT	550	18
blaTEM	TEM-A TEM-B	ATAAAATTCTTGAAGAC TTACCAATGCTTAATCA	1,075	19
blaVEB-1	VEB-FOR VEB-REV	CGACTTCCATTTCCCAGATGC GGACTCTGCAACAAATACGC	642	19
blaSHV-1	SHV-FOR SHV-REV	TGGTTATGCGTTATATTCGCC GCTTAGCGTTGCCAGTGCT	867	20
blaGES-1	GES-1(F) GES-1(R)	ATGCGCTTCATTACGCAC CTATTTGTCCGTGCTCAGG	864	21
blaIMP	IMP-A IMB-B	GAAGGCGTTTATGTTTCATAC GTACGTTTCAAGAGTGATGC	587	22
blaVIM	VIM2004A VIM2004B	GTTTGGTCGCATATCGCAAC AATGCGCAGCACCAGGATAG	382	22
blaKPC	KPC (F) KPC (R)	ATGTCACGTATCGCCGTCT TTACTGCCCCGTTGACGCCCA	880	23
algD	algDF algDR	CGTCTGCCGCGAGATCGGCT- GACCTCGACGGTCTTGCGGA	313	24
lasB	lasBF lasBR	GGAATGAACGAAGCGTTCTCCGAC TTGGCGTCGACGAACACCTCG	284	24
toxA	toxAF toxAR	CTGCGCGGGTCTATGTGCC GATGCTGGACGGGTCGAG-3	270	24
exoS	exoS1-F exoS1-R	TCTGAATTCTCCAGGCGGGTGAACATCA TTAGATCTCACCTGGTATCCAAGGCGA	504	25
exoU	exoU-F exoU-R	GGG AAT ACT TTC CGG GAA GTT- CGA TCT CGC TGC TAA TGT GTT-	428	26

primers, sequences and their product size for detection of extended-spectrum- β -lactamases (ESBLs), metallo- β -lactamases (MBLs) and virulence genes were used according to references as shown in **Table 2** [18-26].

Statistic

Data generated from the study were tabulated as Microsoft Excel sheet and uploaded to Statistical Package for Social Sciences (SPSS version 20). Isolates from clinical sources were calculated for the categorical data by Fisher's exact test, to deter-

mine whether there are any statistical differences in comparison to type of antibiotic resistance and virulence factors. The level of significance was set at a p-value of 0.05 to test the hypothesis of no association; Fisher's exact test is used for small sample size.

Results

A total of 70 *P. aeruginosa* isolates were included in this study, and these were recovered from 40 wound swabs (57.1%), 27 urine samples (38.6%), and 3 pleural fluid samples (4.3%) of patients. Higher rate of isolates were recovered from males 42(60%) than in females 28(40%). Antimicrobial susceptibility testing showed a high to low resistance rates to aztreonam 49 (70%), followed by ceftazidime 32 (45.7%), ciprofloxacin 16 (22.9%), gentamicin 13 (18.6%), piperacillin-tazobactam 11 (15.7%), amikacin 9 (12.9%), and imipenem 6 (8.6%) respectively. All isolates were positive for potential genes of *algD* and *lasB* (100%), followed by *toxA* (90%), *exoS* (34.3%), *exoU* (24.3%) respectively. The rates of detected ESBL genes *bla*TEM, *bla*CTX-m, *bla*SHV-1, *GES-1*, were 3.3%, 6.6%, 10%, 3.3%, and 10% respectively. However, all isolates were negative for *bla*-KPC and *bla*-VIM and IMP genes. The percentages of pigment production were 61.4% for pyocyanin, 37.1% for pyoverdine and 1.4% for pyorubin. (Tables 3, 4, 5 & 6)

Discussion

This study included *P. aeruginosa* isolated from different clinical specimens of patients whom were admitted for medical care at four major hospitals in Khartoum, Sudan. The majority of the isolates were recovered from wound swabs (57.1%), followed from urine (38.6%) and the least percentage was (4.3%) from pleural fluid. The study shows that all isolates have β -hemolytic activity and are mostly

Table 3. Antimicrobial susceptibility of 70 *P. aeruginosa* isolates .

Antimicrobial agent	Resistant Average	
	no	%
Aztreonam	49	70
Ceftazidime	33	47
Ciprofloxacin	16	23
Gentamicin	13	19
Piperacillin- tazobactam	11	16
Amikacin	9	13
Imipenem	6	09

Table 4. MICs of 29 MDR isolates .

Domains/ Facets	MIC ₅₀	MIC ₉₀	MIC Range	Breakpoints for susceptible
	µg/ml	µg/ml	µg/ml)	µg/ml*
Amikacin	7.7	13.9	0.24- >256	≤ 16
Ceftazidime	3.1	5.6	0.75- >256	≤ 8
Imipenem	2.3	4.1	0.38- >32	≤ 2
Ciprofloxacin	2.4	4.2	0.19- > 32	≤ 1

Table 5. Incidence of Virulence genes .

Among 70 <i>P. aeruginosa</i> isolates Gene	<i>P. aeruginosa</i> isolates	
	no	%
AlgD	70	100
LasB	70	100
ToxA	63	90
exoS	24	34.3
exoU	17	24.3

Table 6. Shows distribution of EBLs genes, plasmid-mediated virulence genes of KPC and VIM2/IMP.

GENE	<i>P. aeruginosa</i> isolates	
	no	%
<i>bla</i> TEM	1	3.3
<i>bla</i> CTX-M	2	6.6
<i>bla</i> SHV-1	3	10
<i>bla</i> GES-1	1	3.3
<i>bla</i> VEB	3	10
KPC	0	0
VIM2/IMP	0	0

producing pigmentation, either pyocyanin (61.4%) or pyoverdine (37.1%) and more rarely the pigment pyorubin (1.4%).

Many studies have reported that most important virulence factors of *P. aeruginosa* are production of pigments, including mostly both pyoverdine and pyocyanin and blood hemolysis, which contribute to its pathogenicity during the infection process [27, 28]. However, there are few studies comparing the incidence of pyocyanin and pyoverdine production of *P. aeruginosa* isolates according to their clinical sources. It has been demonstrated that pyocyanin has numerous antagonistic effects and damages host tissues *in vitro* and *in vivo*, including pro-inflammatory and free radical effects [29].

Treatment of infections caused by *P. aeruginosa* has become challenging in recent years due to the ability of this pathogen to rapidly develop resistance to multiple classes of antibiotics. Therefore, the investigation of the genetics of antimicrobial resistance of this organism may help to control and treat its infections by selection of useful antibiotics [30].

This study also showed high to low rates of resistance markers to certain commonly used drugs in treatment of *Pseudomonas* infections including aztreonam (70%), ceftazidime (47%), ciprofloxacin (23%), gentamicin (19%), piperacillin-tazobactam (16%), amikacin (13%) and imipenem (9%) (Table 3). A previous study conducted also in Sudan [31], reported that

P. aeruginosa isolates from clinical samples of patients was 49.3%. The study showed that 74 of the isolates were resistant to carbenicillin [31]. While a new study conducted in Sudan found a multidrug-resistant (MDR) *P. aeruginosa* isolate from urine of patient with a urinary tract infection was highly resistant to many antibiotics including ciprofloxacin, gentamicin, ceftazidime, piperacillin, and meropenem [32].

The most important mechanisms of resistance to carbapenems are production of carbapenemase enzymes, which have the ability to hydrolyze

penicillins, cephalosporins, monobactams and carbapenems. Carbapenem-resistant *Klebsiella* (KPC β -lactamases) hydrolyze most β -lactams, including carbapenems. [33, 34]. Therefore, early detection of this gene will help to avoid incorrect treatment, controlling hospital infections, and preventing the increase spread of resistant strains in hospital and community [35].

The study didn't find any *bla*-*VIM*/IMP and KPC genes in *P. aeruginosa* isolates. Other mechanisms might be responsible for the resistance other than the KPC genes. It is interesting to note here that similar results to ours have been recently reported by a Jordanian study [2].

The study has demonstrated that all *P. aeruginosa* isolates carried potential *AlgD* and *lasB* genes (100%), followed by other toxic genes (Table 6). All MDR isolates were highly associated with these virulence genes compared to less resistant isolates. However, no significant associations were found between antibiotic resistance markers and putative virulence genes neither among MDR nor all *P. aeruginosa* isolates from urine or wound samples *P. aeruginosa* excreted many virulence factors and forms biofilms that contribute to its infection process. Virulence of *P. aeruginosa* is multifactorial, involving both secreted and cell-associated bacterial products such as elastase, alkaline protease, protease IV, exotoxin A and exoenzymeS. However, most studies consider that exotoxin A, exoenzymeS and exoenzymeU are major virulence factors produced by most clinical *P. aeruginosa* strains, and the most toxic effects of these factors are causing host tissue damage [2, 36].

Conclusion

This study demonstrated high rates of antimicrobial resistance markers to most commonly used antibiotics in the treatment of *P. aeruginosa* infections. There was no significant relationship discovered between antimicrobial resistance markers and

potential virulence genes among all *P. aeruginosa* isolates including MDR isolates from either urine or wound samples.

Acknowledgements

The authors are thankful for the Scientific Research Deanship of The University of Jordan, Amman, Jordan for its financial support.

References

1. Shehabi AA, Haider AA, Fayyad MK., Frequency of antimicrobial resistance markers among *Pseudomonas aeruginosa* and *Escherichia coli* isolates from municipal sewage effluent water and patients in Jordan. IAJA 2011;1(1-8).
2. Al Dawodeyah HY, Obeidat N, Abu-Qatouseh LF, Shehab, AA, Antimicrobial resistance and putative virulence genes of *Pseudomonas aeruginosa* isolates from patients with respiratory tract infection. Germs 2018; 8(1), p.31.
3. Paling FP, Wolkewitz M., Depuydt P, De Bus L, Sifakis F, Bonten MJ. et al. 2017. *P. aeruginosa* colonization at ICU admission as a risk factor for developing *P. aeruginosa* ICU pneumonia. Antimicrob Resist Infect Control 2017; 6(1), p.38.
4. Harris AD, Jackson SS, Robinson G, Pineles L, Leekha S, Thom KA, et al. *Pseudomonas aeruginosa* colonization in the intensive care unit: prevalence, risk factors, and clinical outcomes. Infect Cont Hosp Epidemiol 2016; 37(5), 544-548.
5. Driscoll JA, Brody, SL, Kollef M.H. The epidemiology, pathogenesis and treatment of *Pseudomonas aeruginosa* infections. Drugs 2007; 67(3), pp.351-368.
6. Shehabi AA, Masoud H, Balkam Maslamani FA. Common antimicrobial resistance patterns, biotypes and serotypes found among *Pseudomonas aeruginosa* isolates from patient's stools and drinking water sources in Jordan. J Chemother 2005; 17(2), pp.179-183.
7. LaBauve AE, Wargo, MJ. Growth and laboratory maintenance of *Pseudomonas aeruginosa*. Curr Protoc Microbiol 2012; 25(1),6E-1.
8. Persson, A., Study of *Pseudomonas aeruginosa* and different wound dressing products. Master of Science Thesis, Chalmers University Of Technology SE – 412 96 Göteborg Sweden, 2010.
9. Mahmoud AB, Zahran WA, Hindawi GR, Labib AZ, Galal R. . Prevalence of multidrug-resistant *Pseudomonas aeruginosa* in patients with nosocomial infections at a university hospital in Egypt, with special reference to typing methods. J Virol Microbiol 2013; 13, pp.165-59.
10. Rumbaugh KP, Griswold, JA, Hamood AN. *Pseudomonas aeruginosa* strains obtained from patients with tracheal, urinary tract and wound infection: variations in virulence factors and virulence genes. J Hosp Infect 1999; 43(3), 211-218
11. Toder DS, Ferrell SJ, Nezezon JL, Rust L, Iglewski BH. *lasA* and *lasB* genes of *Pseudomonas aeruginosa*: analysis of transcription and gene product activity. Infect Immun 1994; 62(4), 1320-1327.
12. Deretic V, Gill JF, Chakrabarty AM. Gene *algD* coding for GDPmannose dehydrogenase is transcriptionally activated in mucoid *Pseudomonas aeruginosa*. J bacteriol 1987; 169(1), 351-358.
13. Crousilles A, Maunders E, Bartlet S, Fan C, Ukor EF, Abdelhamid Y, et al. Which microbial factors really are important in *Pseudomonas aeruginosa* infections?. Fut Microb 2015; 10(11),1825-1836.
14. Tam VH, Schilling AN, LaRocco MT, Gentry LO, Lolans K, Quinn JP. Prevalence of AmpC over-expression in bloodstream isolates of *Pseudomonas aeruginosa*. Clin Microbiol Infect 2007; 13(4), 413-418.
15. Bonomo RA, Szabo D. Mechanisms of multidrug resistance in *Acinetobacter* species and *Pseudomonas aeruginosa*. Clin infect Dis 2006; 43(2), S49-S56.
16. Clinical Laboratory and Standards Institute (CLSI). Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically; approved standard – tenth edition. CLSI document M07-1. A10 Villanova, PA, USA: CLSI., 2016.
17. Spilker T, Coenye T, Vandamme P, LiPuma JJ. PCR-based assay for differentiation of *Pseudomonas aeruginosa* from other *Pseudomonas* species recovered from cystic fibrosis patients. J Clin Microbiol 2004;42 (5):2074-2079.
18. Pitout JD, Chow BL, Gregson DB, Laupland KB., Elsayed, S. and Church, D.L., 2007. Molecular epidemiology of metallo- β -lactamase-producing *Pseudomonas aeruginosa* in the Calgary Health Region: emergence of VIM2-producing isolates. J Clin Microbiol 2007; 45(2), pp.294-298
19. Jiang X, Ni Y, Jiang Y, Yuan F, Han L, Li M, et al. Outbreak of infection caused by *Enterobacter cloacae* producing the novel VEB-3 beta-lactamase in China. J Clin Microbiol 2000; 43(2):826-31.
20. Fazeli N, Momtaz H. Virulence gene profiles of multidrug-resistant *Pseudomonas aeruginosa* isolated from Iranian hospital infections. Iran Red Crescent Med J. 2014;18; 16 (12).
21. Shacheraghi F, Shakibaie MR, Noveiri H. Molecular Identification of ESBL Genes *bla*GES-1, *bla*VEB-1, *bla*CTX-M, *bla*OXA-1, *bla*OXA-4, *bla*OXA-10 and *bla*PER-1 in *Pseudomonas aeruginosa* Strains Isolated from Burn Patients by PCR, RFLP and Sequencing Techniques. W Acad Sci Eng Technol 2010; 4(1)2010.
22. Pitout JDD, Gregson DB, Poirel L, McClure J-A., Le P, Church DL. Detection of *Pseudomonas aeruginosa* Producing Metallo- β -Lactamases in a Large Centralized Laboratory. J Clin Microbiol 2005; 43(7): 3129–3135.

23. Akpaka PE, Swanston WH, Ihemere HN, Ihemere NH, Correa A, Torres A J. et al. Emergence of KPC-producing *Pseudomonas aeruginosa* in Trinidad and Tobago. J Clin Microbiol 2009;47(8):2670-1.
24. Wolska K, Szewda P. Genetic features of clinical *Pseudomonas aeruginosa* strains. Pol J Microbiol 2009;58:255-60.
25. Shaver CM, Hauser AR. Relative contributions of *Pseudomonas aeruginosa* ExoU, ExoS, and ExoT to virulence in the lung. Infect Immun 2004; 72(12), pp.6969-6977.
26. Mitov I, Strateva T, Markova B. Prevalence of virulence genes among Bulgarian nosocomial and cystic fibrosis isolates of *Pseudomonas aeruginosa*. Braz J Microbiol 2010; 41 (3):588-595.
27. Das T, Manefield M. Pyocyanin promotes extracellular DNA release in *Pseudomonas aeruginosa*. PLoS ONE , 2012; 7, e46718
28. Lau GW, Hassett DJ, Ran H, Kong F. The role of pyocyanin in *Pseudomonas aeruginosa* infection. Tren Mol Med 2004, 10, 599–606.
29. Hall S, McDermott C, Anoopkumar-Dukie, S, McFarland AJ, Forbes A. Perkins AV. Cellular Effects of Pyocyanin, a Secreted Virulence Factor of *Pseudomonas aeruginosa*. Toxins 2016, 8, 236.
30. Lister PD, Wolter DJ, Hanson ND. Antibacterial-Resistant *Pseudomonas aeruginosa*: Clinical Impact and Complex Regulation of Chromosomally Encoded Resistance Mechanisms. Clin Microbiol Rev 2009;22(4):582-610.
31. Abd Elmageed, MD AAE. Molecular Characterization and Sensitivity Patterns of *Pseudomonas aeruginosa* among patients with Different Diseases, PhD Thesis., Sudan University of Sciences and Technology, 2012.
32. Hussain M, Suliman M, Ahmed A, Altayb H, Elneima E. Draft Genome Sequence of a Multidrug-Resistant *Pseudomonas aeruginosa* Strain Isolated from a Patient with a Urinary Tract Infection in Khartoum, Sudan. Genome Announcements 2017;5(16):e00203-17. doi:10.1128/genomeA.00203-17.
33. Castanheira M, Deshpande LM, Costello A, Davies TA, Jones RN. Epidemiology and carbapenem resistance mechanisms of carbapenem-non-susceptible *Pseudomonas aeruginosa* collected during 2009–11 in 14 European and Mediterranean countries. J Antimicrob Chemother 2014;69:1804–1814.
34. Poirel L, Nordmann P, Lagrutta E, Cleary T, and Munoz-Price LS. Emergence of KPC-producing *Pseudomonas aeruginosa* in the United States. Antimicrob Agents Chemother 2010;54:3072.
35. Adler A, Navon-Venezia, S, Moran-Gilad J, Marcos E, Schwartz D, Carmeli Y. Laboratory and clinical evaluation of screening agar plates for detection of carbapenem-resistant Enterobacteriaceae from surveillance rectal swabs. J Clin Microbiol 2011;49: 2239-42.
36. Agnello M, Wong-Beringer A. Differentiation in quinolone resistance by virulence genotype in *Pseudomonas aeruginosa*. PLoS One 2012; 7:e42973

Publish in The International Arabic Journal of Antimicrobial Agents

The Journal is an open access peer-reviewed journal that publishes scientific papers about all aspects of antimicrobials. The journal will publish original research articles, reviews, brief reports and case reports dealing with basic and clinical antibacterial agents, antiviral, antiprotazoals, antituberculous, antifungal and antihelminthes agents. All manuscripts must be prepared in English, and are subject to a rigorous and fair peer-review process. Accepted papers will immediately appear online. The journal aims to advance the knowledge, attitude and the research of chemotherapy in the Arabic world in cooperation with international, national scientific and public societies as well as research centers with similar aims and objectives.