

# Borderline Personality Disorder In Twins: The Preponderance Of Environmental And Genetic Factors

COMMENTARY

Hiroê Alencar Braga<sup>1</sup>,  
Ákila Macêdo Freire<sup>1</sup>,  
Araly Alencar Braga<sup>2</sup>,  
João Marcos Ferreira de  
Lima Silva<sup>1</sup>,  
Modesto Leite Rolim Neto<sup>1</sup>

## Abstract

The borderline personality disorder is a psychiatric disorder that despite being widely researched, still has limitations in addressing certain issues involving the degree of genetic and environmental influence. Although studies indicate the existence of such influences, little had research on different prevalences in twins.

**1** Faculty of Medicine, Estácio-FMJ, Juazeiro do Norte, Ceará, Brazil.

**2** Medical formed by the Federal University of Paraíba –UFPB, PB, Brazil.

## Contact information:

**Hiroê Alencar Braga.**

**Tel:** (055) (88)96054646

**Address:** Rua direita, nº 2,  
Centro, Crato, Ceará, Brazil

✉ [ftphiroe@hotmail.com](mailto:ftphiroe@hotmail.com)

## Keywords

Borderline, Genetic factors, Personality

Borderline personality disorder (BPD) is one of the most studied personality disorders. [1, 2, 3] However, when compared to research on disorders such as depression or other psychiatric disorders, studies on the genetic factors that influence the development of BPD are surprisingly sparse. [3, 4]

BPD, one of the cluster B Axis II personality disorders in DSM-IV [5, 6], is a complex syndrome characterized by pervasive patterns of instability in emotion regulation, interpersonal relationships, self-image and self-control. [6, 7]

It is associated with a number of negative outcomes, including suicidal behavior, frequent emergency room admissions, substance abuse, impaired occupational functioning, and poor quality of interpersonal relationships. [8] Its etiology is not well understood. [9]

Individuals with BPD are well-represented in treatment settings, accounting for 10% of all outpatients and 15-20% of all inpatients [7, 8]. BPD is equally prevalent among men and women and more likely to be diagnosed in early adulthood. [8, 10]

The nine DSM-IV BPD diagnostic criteria specify the core cognitive, behavioral and interpersonal features that identify and differentiate

BPD from other personality and psychiatric disorders. [6] Given that only five of these criteria must be present, there are more than 250 ways to fulfill a BPD diagnosis. It is therefore not surprising that exploratory factor analyses give mixed results, often with three [11, 12, 13] or four [13, 14, 15] highly correlated factors. However, confirmatory factor analyses have generally supported a unidimensional structure. [6, 13, 16, 17, 18]

Initially, research in behavioural and psychiatric genetics focused on disentangling the genetic and environmental influences on traits or disorders. The findings of these studies were highly relevant in showing the influence of genetic factors in the etiology of almost all traits and disorders. Most of these studies assumed that the effects of genes and environment act independently, meaning that the effect of an environmental risk factor does not depend on the genotype. [19]

In twin-family studies, the different degree of genetic relatedness of monozygotic (MZ) and dizygotic (DZ) twin pairs and other first-degree relatives such as siblings is used to identify the relative contribution of genes and environment to the phenotypic variation of a trait. MZ twins share (nearly) all their genes while DZ twins and siblings share on average 50% of their segregating genes. [3, 20]

To date, genetic research on individual differences in BPD has been limited to non-twin family studies and classical twin studies. Family studies have consistently shown increased rates of BPD in family members of BPD patients [8, 21, 22, 23], and twin studies of BPD reported heritability estimates around 40% [8, 24, 25, 26]. Classical twin studies are important to detect whether there are genetic influences on BPD features. [8]

Over a decade ago, "all human behavioral traits are heritable" was stated as the first law of behavior genetics. [27, 28] While provocative at the time, evidence since then has accumulated to suggest heritability estimates of 30% or higher on assessments of cognitive ability, a variety of psychiatric disorders,

and even for most classic personality traits. [28, 29, 30, 31]

Whereas the heritability of common personality traits has been firmly established, the results of the few published studies on personality disorders (PDs) are highly divergent, with some studies finding high heredity and others very low. [32]

Some studies showed that the prevalence of personality disorders varies across studies. [9] In contrast to symptom disorders, few epidemiological studies of personality disorders have been conducted to establish their prevalence. [33]

A problem with assessing personality disorders by means of interview is errors connected with interviewer bias. One reason may be measurement variance related to differences between interviewers. Different interviewers differ in their ability to establish rapport with the twins; they stress different aspects of behavior etc. On the other hand, using the same rater to interview both twins may make the twins too similar of the same reason. A way to overcome the problem is to use self-report questionnaires in addition to interviews. [32]

The currently ascendant theories of BPD etiology are diathesis–stress theories that posit interaction between a child's genetic vulnerability and harsh treatment in the family environment. [34, 35, 36, 37, 38, 39, 40]

Several studies demonstrated that traumatic life events such as sexual and physical abuse, parental divorce or illness or parental psychopathology are important risk factors for the development of BPD. [8, 41, 42, 43, 44] The interaction, however, between the influences of genes and environment on the development of BPD has not been studied. [9]

Recently, several twin and twin family studies provided evidence that genetic factors explain familial clustering of BPD, with heritability estimates ranging from 35% to 45%. [19, 24, 25, 26] However, although many researchers and psychiatrists acknowledge the importance of both traumatic life

events and biological vulnerabilities [8, 19, 45], the joint influence of life events and genetic vulnerability on the development of BPD has not yet been investigated. [19]

There is evidence that inherited and environmental factors each influence BPD etiology. Studies of psychiatric patients show BPD is familial [40, 46] and studies of twin samples show it is heritable. [24, 25, 26, 40, 47]

Some analyses assume that twin resemblance arises from two latent factors: i) additive genes, contributing twice as much to the MZ as to the DZ twin correlation and ii) shared or common environment, which contributes equally to the correlation in MZ and DZ twins. In addition to this common environment, the model also contains individual-specific environment that reflects measurement error and those environmental experiences that make members of a twin pair different. [13]

Only two twin studies so far have provided data on BPD diagnoses and features. Torgersen (1984) [48, 49] reported a monozygotic (MZ) concordance rate of 0.0% and a dizygotic (DZ) concordance rate of 11.1% for BPD, suggesting that shared environmental factors influence the variance in BPD. However, methodological problems of that study limit any conclusions. More recently, Torgersen et al. (2000) [44, 45] reported on the largest twin study to date (n=221 twin pairs) that examined BPD. Results suggested a genetic liability for BPD of 69%, though this heritability estimate must be considered approximate due to the small number of twins, the ascertainment method (sampling those who were treated for mental disorder) and the fact that the zygosity and diagnostic status of co-twins was not hidden from the interviewers. [44]

Has been studied a cohort of twins and our results may not generalize to singletons. However, there is no evidence to suggest that twins and singleton differ in their family psychiatric history, experiences of harsh treatment, or in behavior problems. [40, 46]

There are inherent limitations in all twin studies using phenotypic symptom-based data. The methods used can offer only a coarse approximation of the likely true underlying genetic and environmental structure. For example, the method assumes no gene  $\times$  environment interaction or assortative mating, and it is also dealing with latent liability factors rather than measurable factors. [9]

There are discrepancies in studies of borderline personality disorder in twins as the prevalence of environmental and genetic factors. Future research should focus on possible sources of unique environmental effects and gene environment interaction to develop a comprehensive model of the development of BPD. [3]

## References

1. Blashfield. R. K., & Mcelroy, R. A. (1987). The 1985 journal literature on the personality-disorders. *Comprehensive Psychiatry*. 28. 536-546.
2. Blashfield, R. K., & Intoccia, V. (2000). Growth of the literature on the topic of personality disorders. *American Journal of Psychiatry*, 157, 472-473.
3. Distel MA, Willemsen G, Ligthart L, et al. (2010). Genetic covariance structure of the four main features of borderline personality disorder. *J Pers Disord*. 24(4): 427-444.
4. Crowell, S. E., Beauchaine, T. P., & Linehan, M. M. (2009). A biosocial developmental model of borderline personality; Elaborating and extending Linehan's theory. *Psychological Bulletin*, 135, 495-510.
5. APA (1994). *Diagnostic and Statistical Manual of Mental Disorders*, 4th edn. American Psychiatric Association: Washington, DC.
6. Aggen SH, Neale MC, Røysamb E, Reichborn-Kjennerud T, Kendler KS. (2009). A psychometric evaluation of the DSM-IV borderline personality disorder criteria: age and sex moderation of criterion functioning. *Psychol Med*. 39 (12): 1967-1978.
7. Skodol AE, Gunderson JG, Pfohl B, Widiger TA, Livesley WJ, Siever LJ (2002). The borderline diagnosis I: Psychopathology comorbidity, and personality structure. *Biological Psychiatry* 51, 936-950.
8. Distel MA, Rebollo-Mesa I, Willemsen G, Derom CA, Trull TJ, et al. (2009). Familial Resemblance of Borderline Personality Disorder Features: Genetic or Cultural Transmission? *PLoS ONE* 4(4): e5334.
9. Reichborn-Kjennerud T, Ystrom E, Neale MC, Aggen SH, Mazzeo SE, Knudsen GP, Tambs K, Czajkowski NO, Kendler KS. (2013). Structure of Genetic and Environmental Risk Factors for Symptoms of DSM-IV Borderline Personality Disorder. *Jamapsychiatry*. 70(11): 1206-1214
10. Lenzenweger MF, Lane MC, Loranger AW, Kessler RC (2007). DSM-IV personality disorders in the National Comorbidity Survey Replication. *Biol Psychiat* 62: 553-564.
11. Taylor J, Reeves M. (2007). Structure of borderline personality disorder symptoms in a nonclinical sample. *J Clin Psychol* 63: 805-816.
12. Sanislow CA, Grilo CM, Morey LC et al. (2002). Confirmatory factor analysis of DSM-IV criteria for borderline personality disorder: findings from the collaborative longitudinal personality disorders study. *Am J Psychiatry* 159: 284-290.
13. Kendler KS, Myers J, Reichborn-Kjennerud T. (2011). Borderline personality disorder traits and their relationship with dimensions of normative personality: a web-based cohort and twin study. *Acta Psychiatr Scand*. 123(5): 349-359.
14. Becker DF, Mcglashan TH, Grilo CM. (2006). Exploratory factor analysis of borderline personality disorder criteria in hospitalized adolescents. *Compr Psychiatry* 47: 99-105.
15. De Moor MH, Distel MA, Trull TJ, Boomsma DI. (2009). Assessment of borderline personality features in population samples: is the Personality Assessment Inventory-Borderline Features scale measurement invariant across sex and age? *Psychol Assess* 21: 125-130.
16. Johansen M, Karterud S, Pedersen G, Gude T, Falkum E. (2004). An investigation of the prototype validity of the borderline DSM-IV construct. *Acta Psychiatr Scand* 109: 289-298.
17. Grilo CM, Mcglashan TH, Morey LC et al. (2001). Internal consistency, intercriterion overlap and diagnostic efficiency of criteria sets for DSM-IV schizotypal, borderline, avoidant and obsessive-compulsive personality disorders. *Acta Psychiatr Scand* 104: 264-272.
18. Fossati A, Beauchaine TP, Grazioli F et al. (2006). Confirmatory factor analyses of DSM-IV Cluster C personality disorder criteria. *J Pers Disord* 20: 186-203.
19. Distel MA, Middeldorp CM, Trull TJ, et al. (2011). Life events and borderline personality features: the influence of gene-environment interaction and gene-environment correlation. *Psychol Med*. 41: 849-60.
20. Boomsma, D. I., Busjahn, A., & Peltonen. L. (2002). Classical twin studies and beyond. *Nature Reviews Genetics*, 3, 872-882.
21. Zanarini MC, Frankenburg FR, Yong L, Raviola G, Reich DB, et al. (2004). Borderline psychopathology in the first-degree relatives of borderline and axis II comparison probands. *J Pers Disord* 18: 439-447.
22. White CN, Gunderson JG, Zanarini MC, Hudson JI (2003). Family studies of borderline personality disorder: A review. *Harv Rev Psychiatry* 11: 8-19.
23. Bandelow B, Krause J, Wedekind D, Broocks A, Hajak G, et al. (2005). Early traumatic life events, parental attitudes, family history, and birth risk factors in patients with borderline personality disorder and healthy controls. *Psychiatry Res* 134: 169-179.
24. Distel MA, Trull TJ, Derom CA, et al. (2008). Heritability of borderline personality disorder features is similar across three countries. *Psychol Med*. 38(9): 1219-1229.
25. Kendler KS, Aggen SH, Czajkowski N, Roysamb E, Tambs K, et al. (2008). The Structure of Genetic and Environmental Risk Factors for DSM-IV Personality Disorders A Multivariate Twin Study. *Arch Gen Psychiatry* 65: 1438-1446.
26. Torgersen S, Czajkowski N, Jacobson K, Reichborn-Kjennerud T, Roysamb E, et al. (2008). Dimensional representations of DSM-IV cluster B personality disorders in a population-based sample of norwegian twins: a multivariate study. *Psychol Med* 38: 1617-1625.
27. Turkheimer E (2000). Three laws of behavior genetics and what they mean. *Curr Dir Psychol Sci* 9: 160-164.
28. Lange PAMV, Vinkhuyzen AAE, Posthuma D. (2014). Genetic Influences Are Virtually Absent for Trust. *Plos One* 9 (4): e93880.
29. Bouchard TJ (2004). Genetic influence on human psychological traits. *Curr Dir Psychol Sci* 13: 148-151.

30. Kendler KS (2001). Twin studies of psychiatric illness: An update. *Arch Gen Psychiatry* 58: 1005-1014.
31. Plomin R, Hill L, Craig IW, McGuffin P, Purcell S, et al. (2001). A genome-wide scan of 1842 DNA markers for allelic associations with general cognitive ability: A five stage design using DNA pooling and extreme selected groups. *Behav Genet* 31: 497-509.
32. Torgersen S, Myers J, Reichborn-Kjennerud T, et al. (2012). The heritability of cluster B personality disorders assessed both by personal interview and questionnaire. *J Pers Disord*. 26: 848-66.
33. Torgersen S, Kringlen E, & Cramer V. (2001). The prevalence of personality disorders in a community sample. *Archives of General Psychiatry*, 58, 590-596.
34. Crowell, S. E., Beauchaine, T. P., & Linehan, M. M. (2009). A biosocial developmental model of borderline personality: Elaborating and extending Linehan's theory. *Psychological Bulletin*, 135, 495-510.
35. Fonagy, P., Target, M., & Gergely, G. (2000). Attachment and borderline personality disorder - A theory and some evidence. *Psychiatric Clinics of North America*, 23, 103.
36. Gunderson, J. G., & Lyons-Ruth, K. (2008). BPD's interpersonal hypersensitivity phenotype: A gene-environment-developmental model. *Journal of Personality Disorders*, 22, 22-41.
37. Linehan, M. M. (1993). *Cognitive-behavioral treatment for borderline personality disorder*. New York: Guilford Press.
38. Paris, J. (2005). The development of impulsivity and suicidality in borderline personality disorder. *Development and Psychopathology*, 17, 1091-1104.
39. Zanarini, M. C., & Frankenburg, F. R. (2007). The essential nature of borderline psychopathology. *Journal of Personality Disorders*, 21, 518-535.
40. Daniel WB, Avshalom C, Louise A, et al. (2012). Etiological features of borderline personality related characteristics in a birth cohort of 12-year-old children. *Development and Psychopathology* 24, 251-265
41. Machizawa-Summers S (2007). Childhood trauma and parental bonding among Japanese female patients with borderline personality disorder. *Int J Psychol* 42: 265-273.
42. Zanarini MC, Yong L, Frankenburg FR, Hennen J, Reich DB, et al. (2002). Severity of reported childhood sexual abuse and its relationship to severity of borderline psychopathology and psychosocial impairment among borderline inpatients. *J Nerv Ment Dis* 190: 381-387.
43. Trull TJ (2001). Relationships of borderline features to parental mental illness, childhood abuse, axis I disorder, and current functioning. *J Pers Disord* 15: 19-32.
44. Westen D, Ludolph P, Mislé B, Ruffins S, Block J (1990). Physical and sexual abuse in adolescent girls with borderline personality disorder. *Am J Orthopsychiatry* 60: 55-66.
45. Livesley J (2008). Toward a genetically-informed model of borderline personality disorder. *Journal of Personality Disorders* 22, 42-71.
46. Zanarini, M. C., Barison, L. K., Frankenburg, F. R., Reich, D. B., & Hudson, J. I. (2009). Family history study of the familial coaggregation of borderline personality disorder with Axis I and non-borderline dramatic cluster Axis II disorders. *Journal of Personality Disorders*, 23, 357-369.
47. Torgersen S, Lygren S, Oien PA, Skre I, Onstad S, Edvardsen J, Tambs K, Kringlen E (2000). A twin study of personality disorders. *Comprehensive Psychiatry* 41, 416-425.
48. Torgersen S (1984). Genetic and nosological aspects of schizotypal and borderline personality disorders. A twin study. *Archives of General Psychiatry* 41, 546-554.
49. Distel MA, Trull TJ, Derom CA, et al. (2008). Heritability of borderline personality disorder features is similar across three countries. *Psychol Med*. 38(9): 1219-1229.
50. Johnson, W., Krueger, R. F., Bouchard, T. J., & McGue, M. (2002). The personalities of twins: Just ordinary folks. *Twin Research*, 5, 125-131.

#### Comment on this article:



<http://medicalia.org/>

Where Doctors exchange clinical experiences, review their cases and share clinical knowledge. You can also access lots of medical publications for free. **Join Now!**

**Publish with iMedPub**

<http://www.imed.pub>

International Archives of Medicine is an open access journal publishing articles encompassing all aspects of medical science and clinical practice. IAM is considered a megajournal with independent sections on all areas of medicine. IAM is a really international journal with authors and board members from all around the world. The journal is widely indexed and classified Q1 in category Medicine.