The Efficiency of Electroconvulsive Therapy in the Treatment of Depression in the Elderly

REVIEW

Bruna Furtado Rolim Lima¹, Adriana Araújo de Alencar¹, Daniela Matos Carneiro¹, Inêz Gabrielle Duarte de Sousa¹, Jonathan Almeida Moura¹, Raízza Caroline de Andrade Viana¹, Sonilde Saraiva Januário², Edglê Pedro de Sousa Filho², Uilna Natércia Soares Feitosa², Daniela Cavalcanti e Silva Novais Carvalho², Dayse Christina Rodrigues Pereira Luz², Raimundo Tavares de Luna Neto², Natália Bastos Ferreira², Modesto Leite Rolim Neto^{1,2}

Abstract

The Electroconvulsive Therapy (ECT) is a safe and effective treatment of depression in the elderly. With a high rate of remission may be the treatment of choice in cases of psychotic depression or when antidepressant treatment fails. Although its mechanism of action is unknown the therapeutic effects of ECT may be related to changes in cerebral blood flow. Studies confirm differences in the effects of ECT according to patient age. There are side effects related to cognitive function. ECT is as safe for elderly patients and for young patients.

- 1 Faculty of Medicine, Estácio -FMJ, Juazeiro do Norte, Ceará, Brazil.
- 2 Postgraduate Program in Health Sciences, FMABC, Santo André, São Paulo, Brazil.

Contact information:

Modesto Leite Rolim Neto.

✉ modestorolim@yahoo.com.br

Comment

Major depressive disorder (MDD) in the elderly is a potentially debilitating condition, which rapidly progresses to a severe stage and can be resistant to pharmacological intervention and psychotherapy. [1, 2] In this age group depression is reported to decrease the overall quality of life and because of decreased mobility, compromised food and fluid intake, and a higher incidence of lethal suicide attempts than in younger people, depression should be considered a life-threatening disease. [3, 4]

For the elderly who develop major depression, particularly in cases of psychotic depression or when antidepressant treatment fails, electroconvulsive therapy (ECT) remains a safe and effective treatment with an 80% to 90% remission rate. [5, 6] Although there is little doubt about the efficacy and safety of ECT in clinical practice, the opinions about acceptability, indications and application of ECT differ widely. [7, 8]

Keywords

Electroconvulsive Therapy;
Depression; Elderly.

ECT is a procedure where electrical current is passed through the brain to induce a brief generalized tonic clonic seizure that is therapeutic. This treatment is done under general anesthesia and in a controlled environment where the patient is monitored. [2] Furthermore, the increased blood flow in the thalamus following ECT may also be related to the therapeutic effects of ECT because depression is characterised by symptoms of diencephalic disturbances. [9, 10, 11] Although repeated ECT is generally necessary for ameliorating depressive symptoms, the antidepressant effects of ECT are probably associated with changes in blood flow in the anterior cingulate and medial frontal cortex and thalamus. [11]

The mechanism of action is unknown, but another study has shown a significant increase in cerebral gabaaminobutyric acid (GABA) after ECT. [2, 12] PET scans done pre-ECT and after a course of ECT showed a decrease in the cerebral metabolism in prefrontal and parietal areas, and this reduction correlated with improvement in depression. [2, 13]

Middle-aged and elderly patients have a higher remission rate after a course of ECT than younger patients. [14, 15, 16, 17] But, the association between age and ECT outcome could be dependent on the propensity of the elderly to suffer from certain types of depression. [18, 19] The efficacy of ECT was significantly superior in patients with psychotic depression. [20] Moreover, electroconvulsive therapy deserves a more prominent position in the treatment of elderly patients with severe unipolar depression. [4]

A number of case studies report efficacy of ECT in demented patients, patients with cerebrovascular disorders and depressed patients with Parkinson's disease. [8] Furthermore, ECT was well tolerated by the elderly even in patients older than 80 years. [21]

Several studies have investigated the capacity of depressed patients to consent to treatment or research. [6, 22, 23] Those who were older had more

difficulty in fully understanding all the relevant information concerning ECT. [6]

When studying cognitive side effects of ECT, it is important to take into account the most relevant treatment and clinical characteristics that may influence these cognitive side effects, such as electrode placement, age, and the severity of depressive symptoms. [24] There is convincing evidence from studies of patients of mixed ages that bilateral (usually bitemporal) ECT results in more cognitive side-effects than unilateral treatment. [25] But, in general, older patients may be more vulnerable to cognitive problems than younger patients because age in itself may lead to cognitive frailty, but also because of comorbid conditions, such as diabetes, hypertension, cerebro-vascular disease, dementia and M. Parkinson. [26]

The manner via which ECT should be administered to optimize the relationship between efficacy and negative cognitive effects remains controversial. [27, 28] Nevertheless, multiple studies have supported the safety and efficacy of ECT in the elderly, even in patients older than 75 years with multiple medical comorbidities and cognitive impairment. [14, 15, 29] One study mentioned the possibility of developing post-ECT delirium or dementia in older patients although these are not known to be side effects of ECT. [21]

ECT improved speed of information processing, mental control, memory (all aspects) and perception (visual organization), recovery of depression was especially associated with improvement of memory (verbal learning), higher speed of information processing and improvement of executive functioning. [26, 30]

When ECT was compared with antidepressants it was almost always reported to have superior therapeutic efficacy. [21] By using more specific and sensitive neuropsychological instruments we should be able to detect cognitive dysfunction related to independency in daily living after ECT. [26]

ECT is the most effective treatment in severely depressed elderly patients and can have manageable side effects, with transient adverse events and limited mortality using modern evidence-based protocols. [8, 14, 31] For those with a limited understanding of the treatment being recommended, educational interventions can be helpful. [6]

Yet, up to 20% of elderly patients may not respond to ECT, and the elderly are more susceptible to the cognitive side effects, including delirium and acute cardiovascular and other medical complications associated with ECT. [14] Age-associated structural brain changes may be associated with poorer response to ECT. [19, 32]

Alternative therapies are clearly needed for depressed geriatric patients who are resistant or intolerant to available somatic therapies. [14] Electroconvulsive therapy (ECT) is an important treatment option in late life depression [33, 34] although it was first introduced in 1938 as a treatment for schizophrenia. [11] A large number of well-designed studies support the use of drugs and ECT for the safe and effective treatment of depressed elderly individuals. [21] Importantly, this comparability in outcome across age groups was also evident 1-3 years after ECT. As the level of adverse events was not increased after ECT for subjects in any age group, we conclude that ECT is as effective and safe for older depressed patients as it is for younger ones. [19].

References

1. Alexopoulos GS, Meyers BS, Young RC, Kakuma T, Feder M, Einhorn A, Rosendahl E. Recovery in geriatric depression. *Arch Gen Psychiatry*. 1996; 53: 305-12.
2. Manepalli J, Sapkota N. Neuromodulation Therapies in the Elderly Depressed Patient. *Geriatric Psychiatry*. 2014; 3: 229-236
3. Baldwin RC. Poor prognosis of depression in elderly people: causes and actions. *Ann Med* 2000; 32: 252-6.
4. Spaans HP, Sienaert P, Bouckaert F, Van den Berg JF, Verwijk E, Kho KH, Stek ML, Kok RM. Speed of remission in elderly patients with depression: electroconvulsive therapy v. medication. *The British Journal of Psychiatry* 2015; 206: 67-71.
5. Greenberg L, Fink M. The use of electroconvulsive therapy in geriatric patients. *Clin Geriatr Med* 1992; 8: 349-354.
6. Lapid MI, Rummans TA, Pankratz VS, Appelbaum PS. Decisional Capacity of Depressed Elderly to Consent to Electroconvulsive Therapy. *J Geriatr Psychiatry Neurol* 2004; 17: 42-46
7. Philpot M, Treloar A, Gormley N, Gustafson L. 2002. Barriers to the use of electroconvulsive therapy in the elderly: a European survey. *Eur Psych* 17: 41-45.
8. Van der Wurff FB, Stek ML, Hoogendijk WJG, Beekman ATF. The efficacy and safety of ECT in depressed older adults, a literature review. *Int J Geriatr Psychiatry* 2003; 18: 894-904.
9. N - Carney MWP, Sheffield BF. Electroconvulsive therapy and diencephalon. *Lancet*, i, 1973; 1505-1506.
10. Abrams R, Taylor MA. Diencephalic stimulation and the effects of ECT in endogenous depression. *British Journal of Psychiatry* 1976; 129: 482-485.
11. Takano H, Motohashi N, Uema T, Ogawa K, Ohnishi T, Nishikawa M, Kashima H, Matsuda H. Changes in regional cerebral blood flow during acute electroconvulsive therapy in patients with depression. *British Journal Of Psychiatry* 2007; 190: 63-68.
12. Sanacora G, Mason GF, Rothman DL, et al. Increased cortical GABA concentration in depressed patients receiving ECT. *Am J Psychiatry*. 2003; 160(3): 577-9.
13. Nobler MS, Oquendo MA, Kegeles LS, Malone KM, Campbell CC, Sackeim HA, Mann JJ. Decreased regional brain metabolism after ECT. *Am J Psychiatry*. 2001; 158: 305-8.
14. Riva-Posse P, Hermida AP, McDonald WM. The Role of Electroconvulsive and Neuromodulation Therapies in the Treatment of Geriatric Depression. *Psychiatr Clin N Am* 36; 2013; 607-630.
15. Tew JD Jr, Mulsant BH, Haskett RF, et al. Acute efficacy of ECT in the treatment of major depression in the old-old. *Am J Psychiatry* 1999;156(12): 1865-70.
16. Flint AJ, Gagnon N. Effective use of electroconvulsive therapy in late-life depression. *Can J Psychiatry* 2002; 47(8): 734-41.
17. Greenberg RM, Kellner CH. Electroconvulsive therapy: a selected review. *Am J Geriatr Psychiatry* 2005; 13(4): 268-81.

18. Benbow SM. The role of electroconvulsive therapy in the treatment of depressive illness in old age. *Br. J. Psychiatry* 1989; 155: 147-152.
19. Brodaty H, Hickie I, Mason C, Prenter L. A prospective follow-up study of ECT outcome in older depressed patients. *Journal of Affective Disorders* 2000; 60: 101-111.
20. Birkenhäger TK, Pluijms EM, Ju MR, Mulder PG, Van den Broek WW. Influence of age on the efficacy of electroconvulsive therapy in major depression: A retrospective study. *Journal of Affective Disorders* 2010; 126: 257-261.
21. W - Salzman C, Wong E, Wright BC. Drug and ECT Treatment of Depression in the Elderly, 1996-2001: A Literature Review. *Biol Psychiatry* 2002; 52: 265-284.
22. Appelbaum PS, Grisso T, Frank E, et al. Competence of depressed patients for consent to research. *Am J Psychiatry* 1999; 156: 1380-1384
23. Roth LH, Lidz CW, Meisel A, et al. Competency to decide about treatment or research. *Int J Law Psychiatry* 1982; 5: 29-50.
24. Verwijk E, Comijs HC, Kok RM, Spaans HP, Tielkes CEM, Scherder EJA, Stek ML. Short- and long-term neurocognitive functioning after electroconvulsive therapy in depressed elderly: a prospective naturalistic study. *International Psychogeriatrics* 2014; 26: 2, 315-324.
25. O'Connor DW, Gardner B, Eppingstall B, Tofler D. Cognition in elderly patients receiving unilateral and bilateral electroconvulsive therapy: A prospective, naturalistic comparison. *Journal of Affective Disorder* 2010; 124: 235-240.
26. Tielkes CEM, Comijs HC, Verwijk E, Stek ML. The effects of ECT on cognitive functioning in the elderly: a review. *Int J Geriatr Psychiatry* 2008; 23: 789-795.
27. Bjølseth TM, Engedal K, Benth JS, Dybedal GS, Gaarden TL, Tanum L. Clinical efficacy of formula-based bifrontal versus right unilateral electroconvulsive therapy (ECT) in the treatment of major depression among elderly patients: A pragmatic, randomized, assessor-blinded, controlled trial. *Journal of Affective Disorders* 2015; 175: 8-17.
28. Sackeim HA, Prudic J, Devanand DP, Nobler MS, Lisanby SH, Peyser S, Fitzsimons L, Moody BJ, Clark J. A prospective randomised double-blind comparison of bilateral and right unilateral electroconvulsive therapy at different stimulus intensities. *Arch.Gen.Psychiatry* 2000; 57: 425-434.
29. Cattan RA, Barry PP, Mead G, et al. Electroconvulsive therapy in octogenarians. *J Am Geriatr Soc* 1990; 38(7): 753-8.
30. Bosboom PR, Deijen JB. Age-related cognitive effects of ECT and ECT-induced mood improvements in depressive patients. *Depress Anxiety* 2006; 23: 93-101.
31. Damm J, Eser D, Schule C, et al. Influence of age on effectiveness and tolerability of electroconvulsive therapy. *J ECT* 2010; 26(4): 282-8.
32. Hickie I, Scott E, Wilhelm K, Brodaty H. Subcortical hyperintensities on magnetic resonance imaging in patients with severe depression – a longitudinal evaluation. *Biol. Psychiatry* 1997; 42: 367-374.
33. UK ECT Review Group. Efficacy and safety of ECT in depressive disorders: a systematic review and meta-analysis 2003; 361: 799-808.
34. Stek ML, van der Wurff FB, Uitdehaag BMJ, Beekman ATF, Hoogendijk WJG. ECT in the treatment of depressed elderly: lessons from a terminated clinical trial. *Int. J. Geriatr. Psychiatry* 2007; 22: 1052-1054.

Comment on this article:

<http://medicalia.org/>

Where Doctors exchange clinical experiences, review their cases and share clinical knowledge. You can also access lots of medical publications for free. **Join Now!**

Publish with iMedPub

<http://www.imed.pub>

International Archives of Medicine is an open access journal publishing articles encompassing all aspects of medical science and clinical practice. IAM is considered a megajournal with independent sections on all areas of medicine. IAM is a really international journal with authors and board members from all around the world. The journal is widely indexed and classified Q1 in category Medicine.