Prevention of Prostate Cancer in Primary Care: an Integrative Review

REVIEW

Laryssa Lins de Araújo¹, Thiago Lívio Barbosa Pereira²,
Marilena Maria de Souza³,
Renata Lívia Silva Fonseca Moreira de Medeiros⁴,
Janaina Ferreira Moreira⁵, Paula Frassinetti Oliveira Cezário¹,
Arielle Wignna Brasil Abrantes⁶,
Gabrielle Moraes Arruda Costa², Jesana Sá Damasceno Moraes⁷,
Erinalda Araújo Ramos⁸, Kátia Alves Soares⁹,
Francisca Cristiane Pessoa Lima¹⁰, Francisca Karla Esmeraldo Neves¹¹,
Janaíne Chiara Oliveira Moraes¹², Renata Maia de Medeiros¹³

- 1 Specialist in Family Health, Federal University of Paraíba. João Pessoa*.
 - 2 Bachelor of Nursing, Federal University of Campina Grande. Cajazeiras*.
 - 3 Doctor in Medicine and Health, Federal University of Bahia. Salvador**.
 - 4 Master of the Nursing Postgraduation Program, Federal University of Paraíba. João Pessoa*.
 - 5 Specialist in Intensive Therapy, Federal University of Paraíba. João Pessoa*.
 - 6 Specialist in Nursing Intensive Care, Federal University of Campina Grande. Cajazeiras*.
 - 7 Specialist in Health Management, State University of Paraíba. Cajazeiras*.
 - 8 Nursing Student, College Santa Maria. Cajazeiras*.
 - 9 Bachelor of Nursing, College Santa Maria. Cajazeiras*.
 - 10 Specialist in Urgency and Emergency, Occupational Health and Occupational Health Nursing, Cajazeiras*.
 - 11 Specialist, Regional Hospital of Cajazeiras. Cajazeiras*.
 - 12 Academic Unit of Nursing, Teacher Training Center, Federal University of Campina Grande. Cajazeiras*.
 - 13 Nursing Student, Federal University of Paraíba. João Pessoa*.
- * Paraíba, Brazil. ** Bahia, Brazil.

Abstract

Introduction: Prostate cancer is one of the most prevalent diseases in the male population worldwide. This type of cancer can be detected early in a basic health unit through the evaluation of the prostate with the digital rectal exam, and the serum prostate specific antigen.

Objective: Evaluating the prevention of prostate cancer in primary care from the analysis of scientific productions.

Method: It was decided to carry out an integrative review, using the descriptors: human health, primary care, prostate, prostate cancer and men's health; the databases consulted were LILACS, SciELO and PubMed, from 2008 to September 2015.

Results: The findings showed that interventions associated to health education may result in changes of behavior and habits of health, particularly for individuals who have risk factors or are in the early stages of the disease.

Conclusion: it is expected that this study will encourage professionals of the units to initiate changes that result in a better acceptance of men to action and better efficiency indexes in the collection and

Contact information:

Thiago Lívio Barbosa Pereira.

✉ thigolivio7@hotmail.com

treatment of patients with cancer. In addition, it is necessary that these actions advance and become routine in institutions, not restricted to the campaigns of "Blue November", but forming part of the daily planning of the Family Health Strategy.

Keywords

Men's Health; Cancer of the Prostate; Prevention; Community Health Services.

Introduction

The National Integral Attention to Men's Health Policy (NIAMHP) was established in August 2009 under the Ordinance 1944 and established by the Unified Health System (UHS). This policy has the purpose to demystify the idea of invulnerability on male health, provide health professionals a greater understanding about the disease process of these individuals, and develop actions and strategies that encourage men to attend health facilities of primary care [1].

Most men do not seek primary care services for the prevention and treatment of pathological processes in the early stages, preferring to use the specialized care services of urgent moments or when the disease is installed, thus causing the increase in morbidity and mortality rates in this population and the increased costs to the public purse. Therefore, the objective of NIAMHP strengthen and qualify the primary care so that health care is not just restricted to recovery, ensuring, above all, health promotion and prevention to preventable diseases [1].

Among the diseases that most affect the male population, prostate cancer deserves attention because it is considered the sixth most common type of cancer worldwide and the second most prevalent in men, accounting for about 10% of all malignancies. In 2014, we calculated the emergence of 68.800 new cases, representing an estimated risk of 70.42 new cases per 100.000 men [2].

This cancer can be detected early and simply in a Family Health Strategy (FHS), through the evaluation of the prostate with the digital rectal exams and the serum levels of Prostate Specific Antigen (PSA) [3].

From the studies conducted on human health, the experience in health facilities and the itinerant design of the Cajazeiras Regional Hospital, it was noted the demand of this population for actions that address the prevention of diseases, especially prostate cancer. Therefore, we opted for this study because of the need of passing on information that may contribute to a better assistance to the male population in the FHS.

Thus, the objective of this study is to assessing the prevention of prostate cancer in primary care from the analysis of scientific productions.

Method

We opted for the realization of an integrative literature review, because it addresses, in a satisfactory manner, the established objective, in addition to performing as a broad method, which allows the incorporation of several studies, thus resulting in a more comprehensive knowledge about the subject studied [4].

The integrative review process includes six distinct stages, similar to the stages of performing a standard search:

1st) Issue identification and selection of the research question. This step begins with the definition of the problem and formulates a research hypothesis, which must have relevance to the health of individuals. In this sense, the guiding question of the study was: How is taking place prevention of prostate cancer in primary care?

2nd) Criteria for the selection of the sample. on the 1st of October 2015 there was conducted to

search the publications by searching in the databases of the Virtual Health Library (VHL): SciELO (Scientific Electronic Library Online), LILACS (Latin American and Caribbean Health Sciences) and international database PubMed (Medical Published - serve of the US National Library of Medicine). They used the descriptors: human health, prostate cancer, primary care, men's health, prostate cancer and Family Health. Inclusion criteria articles were: scientific works published in Portuguese and English, with abstracts and full text available free of charge in full in the period between 2008 and September 2015. In the first search performed using the above descriptors, found 763 articles, and 316 in Portuguese and 447 in English.

3rd) Identification of pre-selected and selected studies. First held up reading the summaries of each article, being selected 21 publications in PubMed, 15 and 6 in the SciELO and LILACS. Subsequently, the complete reading of the pre-selected articles and verified their compliance with the inclusion criteria and the guiding question of the survey was conducted, which resulted in the selection of 13 articles, which were 9 and 4 of the SciELO and LILACS.

4th) Categorization of studies. This step is considered similar to the collection of conventional data, for which we used a tool [5] that allows the researcher to evaluate, individually, selected studies, analyzing them methodologically and in relation to synthesis of the findings. Thus, studies were listed through a careful reading and inclusion criteria chosen.

5th) Analysis and interpretation of results. For analysis and subsequent synthesis of the articles there was prepared a framework containing the following aspects relevant to the selected theme: author names, article title, journal and publication year. Then, the results were discussed according to the relevant literature.

6th) Presentation of synthesis of knowledge. The knowledge gained by conducting this review was presented as a document, which forms the explicit content in this scientific work.

Results

Selected publications-specifications

Table 1 presents the authors, titles, periodicals and years relating to each publication analyzed. Journals that published more articles were: Brazilian Journal

Table 1. Characterization of publications according to the authors, title, journal and year of publication, in the period from 2008 to 2015.

Authors	Title	Journal, Year
Medeiros AP, Menezes MFB, Napoleão AA [3]	Risk factors and prevention of prostate cancer: subsidies for nursing.	Brazilian Journal of Nursing, 2011
Gomes R et al. [6]	The scratching of masculinity: a discussion on the DRE as prostate cancer prevention.	Science & Public Health, 2008
Gonçalves IR, Padovani C, Popim RC [7]	Epidemiological and demographic characteristics of men with prostate cancer.	Science & Public Health, 2008
Gomes R et al. [8]	Prevention of prostate cancer: a literature review.	Science & Public Health, 2008
Seoane AF, Fortes PAC [9]	The perception of the user of the family health program about the privacy and confidentiality of his information.	Health and Society, 2009
Paiva EP, Motta MCS, Griep RH [10]	Knowledge, attitudes and practices regarding the detection of prostate cancer.	Acta Paulista of Nursing, 2010
Migowski A, Silva GA [11]	Survival and prognostic factors of patients with clinically localized prostate cancer.	Journal of Public Health, 2010

Authors	Title	Journal, Year
Amorim VMSL et al. [12]	Factors associated with the screening tests for prostate cancer: A population-based study.	Notebook of Public Health, 2011
Souza LM, Silva MP, Pinheiro IS [13]	A touch on masculinity: the prevention of prostate cancer in traditionalist gauchos.	Gaucha Journal of Nursing, 2013
Belinelo RGS et al. [14]	Screening tests for prostate cancer: experience of men.	Anna Nery School, 2014
Nogueira HL, Neves JB [15]	Prevention of prostate cancer: role of nurses in primary health care units.	Integrated Nursing Magazine - Ipatinga, 2013
Santos ACF, Silva EM, Pacheco JBP[16]	Evaluation of the first campaign for the diagnosis and prevention of prostate cancer promoted by the municipal government of Anapolis-GO.	Yearbook of Undergraduate Research Student Production, 2010
Silva NA et al.[17]	Man Health promotion in primary health care services.	in Extension, 2014

Source: Prepared by the authors. Database SciELO and LILACS (2008-2015).

of Nursing [3], Science & Public Health [6-8], Health and Society [9], Acta Paulista of Nursing [10], Journal of Public Health [11], Notebook of Public Health [12], Gaucha Journal of Nursing [13], Anna Nery School [14], Integrated Nursing Magazine - Ipatinga [15], Yearbook of Undergraduate Research Student Production [16], in Extension [17].

Discussion

The FHS was developed as a plan to guide the organization of the health system and respond to the population's needs, requiring an understanding of health as a social right and favors the facing of social determinants to promote it. Thus, the proper organization of the services available in the FHS contribute to the assistance provided under primary care, with positive impacts on health population [18].

According to Ordinance 648 from March 28th, 2006 [19], all FHS's team member professionals have as common assignments health promotion, disease prevention, treatment and monitoring of diseases, and activities permanent health education. From this, the creation of the FHS provided a focused attention on the family, understood and perceived by their physical and social environment, thus enabling the FHS team to work together with these persons

a broader understanding of the health/disease process and the need to interventions that transcend healing practices [9, 15, 20].

Thus, arises the need to address human health and all the complexities involving the care of these individuals. To do this, there was set up to NIAMHP, which directed the focus of attention of professionals to the most common disorders in the male population, especially in relation to prostate cancer. This disease is the sixth most common malignancy in the world in number of new cases, the third most common cancer in the male population and more neoplasia presented by European men, American and some parts of Africa [3, 6, 8, 10].

In recent years there have seen a significant increase in incidence rates of this cancer in Brazil, which can be explained by the increase in population life expectancy, the constant campaigns to identify the disease, the development of diagnostic methods and quality systems Brazilian information, in addition to environmental and dietary influences, such as high energy consumption, intake of red meat, fats and milk [14, 21-22].

The study of prostate cancer is of great importance to public health, due to its high incidence and high clinical cure rates in these patients when the disease is detected in early stages [22-23]. The tests most commonly used to screen cancer are the

digital rectal examination and serum PSA. The first has limitations when the prostate has a not palpable lesion, common in early stages, and when changed, the differentiation between malignant and benign lesion is quite complex. The second examination, serum PSA in clinical use since 1986, is the tumor marker most often used for screening for prostate cancer and plays an important role in the early diagnosis and prevention of complications of the disease, direct impact in reducing of morbidity and mortality [6, 12-13, 24].

The prevention of prostate cancer should be performed primarily in primary care services such as FHS, and the role of professionals receiving the male population in a universal manner, comprehensive and equitable, given the advocated by UHS principles and guidelines, and promoting increased access to the right information on health, especially on prostate cancer, contributing thus to face the risk factors [11, 13, 15]. Moreover, it should also be encouraged by the FHS professionals, adopting healthy eating habits and physical activity as a means of prevention against prostate cancer and other common injuries to this population [16-17].

In Brazil, about 60% to 70% of cases are diagnosed when the disease is already widespread, given that, in most cases, this type of cancer has a very slow growth period, taking years to the individual presents any symptom [7, 17]. We observe, then, the great need to target the male population about the issues surrounding prostate cancer and its prevention and early detection in primary care [16].

Through this context notes the importance of the FHS for the prevention of prostate cancer and other diseases prevalent in the male population and society in general, working efficiently and resolute manner, and reducing morbidity and mortality from preventable causes [9, 15].

The FHS brings in its conception a redesign of care model that emphasizes the link with care lines of logic in which every citizen has the right to be ca-

red for by a team of full and resolute manner, with therapeutic projects caregivers solidarity with the demands and/or the suffering of users, also being co-responsibilities of producers and autonomy or self-government on the part of those involved in this process [25].

Conclusion

By analyzing the studies included in the research, it was observed that the strategies used for the prevention of prostate cancer in the context of primary health care, are the examination of digital rectal examination and serum PSA. In addition, some also mention encouraging the adoption of lifestyles and healthy food, contributing to the prevention of numerous injuries to men's health.

Searching for information in the scientific literature and pass on knowledge to the people is an extremely important action for public health, allowing the exchange of knowledge and experiences between professionals and users, thus contributing to the formation and/or strengthening of the bond. So, through health education and the commitment of the FHS teams, strategies can be developed to encourage men to seek more frequently the primary care services for health promotion and prevention of diseases, not limited only to monitoring and treatment health problems.

It is hoped that this study will encourage practitioners to start on changes that result in better acceptance of men to action and best efficiency ratios in the collection and treatment of patients with cancer. It is hoped, too, that these actions advance and become routine in health institutions as part of the daily planning of the FHS.

References

1. Ministério da Saúde (BR). *Política Nacional de Atenção Integral à Saúde do Homem (Princípios e Diretrizes)* [Internet]. Brasília: Ministério da Saúde; 2008. Available from: http://bvsm.s.saude.gov.br/bvs/publicacoes/politica_nacional_atencao_homem.pdf

2. Instituto Nacional do Câncer (INCA). Ministério da Saúde. *Programa Nacional de Controle do câncer da Próstata: Documento de Consenso* [Internet]. Rio de Janeiro: INCA; 2002. Available from: http://bvsmms.saude.gov.br/bvs/publicacoes/cancer_da_prostata.pdf
3. Medeiros AP, Menezes MFB, Napoleão AA. Fatores de risco e medidas de prevenção do câncer de próstata: subsídios para a enfermagem. *Rev Bras Enferm* [Internet], Brasília, 2011 Mar/Apr; 64(2): 385-8. DOI: <http://dx.doi.org/10.1590/S0034-71672011000200027>
4. Mendes KDS, Silveira RCCP, Galvão CM. Revisão integrativa: método de pesquisa para a incorporação de evidências na saúde e na enfermagem. *Texto Contexto Enfermagem* [Internet], 2008; 17(4): 758. DOI: <http://dx.doi.org/10.1590/S0104-07072008000400018>
5. Ursi SE, Galvão, CM. Perioperative prevention of skin injury: an integrative literature review. *Rev Latino-Am Enfermagem* [Internet], Ribeirão Preto 2006 Jan/Feb; 14(10): 124-31. DOI: <http://dx.doi.org/10.1590/S0104-11692006000100017>
6. Gomes R, Nascimento EF, Rebello LEFS, Araújo FC. As arranhaduras da masculinidade: uma discussão sobre o toque retal como medida de prevenção do câncer prostático. *Ciênc Saúde Coletiva* [Internet], Rio de Janeiro, 2008 Nov/Dec; 13(6): 1975-84. DOI: <http://dx.doi.org/10.1590/S1413-81232008000600033>
7. Gonçalves IR, Padovani C, Popim RC. Caracterização epidemiológica e demográfica de homens com câncer de próstata. *Ciênc Saúde Coletiva* [Internet], Rio de Janeiro, 2008 July/Aug; 13(4): 1337-42. DOI: <http://dx.doi.org/10.1590/S1413-81232008000400031>
8. Gomes R, Rebello LEFS, Araújo FC, Nascimento EF. A prevenção do câncer de próstata: uma revisão da literatura. *Ciênc Saúde Coletiva* [Internet], Rio de Janeiro, 2008 Jan/Feb; 13(1): 235-46. DOI: <http://dx.doi.org/10.1590/S1413-81232008000100027>
9. Seoane AF, Fortes PAC. A percepção do usuário do programa saúde da família sobre a privacidade e a confidencialidade de suas informações. *Saúde Soc* [Internet], São Paulo, 2009 Jan/Mar; 18(1): 42-9. DOI: <http://dx.doi.org/10.1590/S0104-12902009000100005>
10. Paiva EP, Motta MCS, Griep RH. Conhecimentos, atitudes e práticas acerca da detecção do câncer de próstata. *Acta Paul Enferm* [Internet], 2010; 23(1): 88-93. Available from: <http://www.scielo.br/pdf/apv/v23n1/14.pdf>
11. Migowski A, Silva GA. Sobrevida e fatores prognósticos de pacientes com câncer de próstata clinicamente localizado. *Rev Saúde Pública* [Internet], São Paulo, 2010 Apr; 44(2): 344-52. DOI: <http://dx.doi.org/10.1590/S0034-89102010000200016>
12. Amorin VMSL, Barros MBA, César CLG, Goldbaum M, Carandina L, Alves MCGP. Fatores associados à realização dos exames de rastreamento para o câncer de próstata: um estudo de base populacional. *Cad Saúde Pública* [Internet], Rio de Janeiro, 2011 Feb; 27(2): 347-56. DOI: <http://dx.doi.org/10.1590/S0102-311X2011000200016>
13. Souza LM, Silva MP, Pinheiro IS. Um toque na masculinidade: a prevenção do câncer de próstata em gaúchos tradicionalistas. *Rev Gaúcha Enferm* [Internet], Porto Alegre, 2011 Mar; 32(1): 151-8. DOI: <http://dx.doi.org/10.1590/S1983-14472011000100020>
14. Belinelo RGS, Almeida SM, Oliveira PP, Onofre PSC, Viegas SMF, Rodrigues AB. Exames de rastreamento para o câncer de próstata: vivência de homens. *Esc Anna Nery* [Internet], Rio de Janeiro, 2014 Oct/Dec; 18(4): 697-704. DOI: <http://dx.doi.org/10.5935/1414-8145.20140099>
15. Nogueira HL, Neves JB. Prevenção do câncer da próstata: atuação dos enfermeiros nas unidades de atenção primária a saúde. *Revista Enfermagem Integrada – Ipatinga: Unileste* [Internet] 2013 Jul/Ago; 6(1): 1098-1109. Available from: <http://www.unilestemg.br/enfermagemintegrada/artigo/v6/07-prevencao-do-cancer-da-prostata-atuacao-dos-enfermeiros-nas-unidades-de-atencao.pdf>
16. Santos ACF, Silva EM, Pacheco JBP. Avaliação da primeira campanha para o diagnóstico e prevenção do câncer de próstata promovido pela prefeitura municipal de Anápolis-GO. *Anuário da Produção de Iniciação Científica Discente* [Internet], 2010; 13(17): 25-42. Available from: <http://saic.anhanguera.com/index.php/PAP/article/view/1838>
17. Silva NA, Dias MP, Silva DA, Dias LP. Promoção da saúde do homem nos serviços de atenção primária à saúde. *Em Extensão* [Internet], Uberlândia, 2014; Jan/Jul; 13(1): 82-8. Available from: <http://www.seer.ufu.br/index.php/revextensao/article/view/23996/14683>
18. Giovanella L, Mendonça MHM, Almeida PF, Escorel S, Senna MCM, Fausto MCR et al. Saúde da família: limites e possibilidades para uma abordagem integral de atenção primária à saúde no Brasil. *Ciênc Saúde Coletiva* [Internet], Rio de Janeiro, 2009 May/June; 14(3): 783-94. DOI: <http://dx.doi.org/10.1590/S1413-81232009000300014>
19. Ministério da Saúde (BR). *Portaria Nº 648/GM de 28 de março de 2006* [Internet]. Aprova a Política Nacional de Atenção Básica, estabelecendo a revisão de diretrizes e normas para a organização da Atenção Básica para o Programa Saúde da Família (PSF) e o Programa Agentes Comunitários de Saúde (PACS). Available from: http://bvsmms.saude.gov.br/bvs/publicacoes/prtgM648_20060328.pdf

20. Santos KT, Saliba NA, Moimaz SAS, Arcieri RM, Carvalho ML. Agente comunitário de saúde: perfil adequado a realidade do Programa Saúde da Família? *Ciênc Saúde Coletiva* [Internet], Rio de Janeiro, 2011; 16(Supl. 1): 1023-8. DOI: <http://dx.doi.org/10.1590/S1413-81232011000700035>
21. Instituto Nacional do Câncer (INCA). Ministério da Saúde. *Câncer de Próstata* [Internet]. 2014. Available from: <http://www2.inca.gov.br/wps/wcm/connect/tiposdecancer/site/home/prostata/definicao>
22. Srougi M, Ribeiro LA, Piovesan AC, Colombo JR, Nesrallah A. Doenças da próstata. *Rev Med* [Internet], São Paulo, 2008 Jul/Set; 87(3): 166-77. Available from: http://fm.usp.br/gdc/docs/revistadc_166_07-doencasdaprostata.pdf
23. Jemal A, Siegel R, Ward E, Murray T, Xu J, thun MJ. Cancer statistics, 2007. *CA Cancer J Clin* [Internet], 2007 Jan/Feb; 57(1): 43-66. PMID: 17237035.
24. Castro HAS, IARED W, Shigueoka DC, Morão JE, Ajzen S. Contribuição da densidade do PSA para predizer o câncer da próstata em pacientes com valores de PSA entre 2,6 e 10,0 ng/ml. *Radiol Bras* [Internet], São Paulo, 2011 Jul/Ago; 44(4): 205-9. DOI: <http://dx.doi.org/10.1590/S0100-39842011000400003>
25. Schwartz TD, Ferreira JTB, Maciel ELN, Lima RCD. Estratégia Saúde da Família: avaliando o acesso ao SUS a partir da percepção dos usuários da Unidade de Saúde de Resistência, na região de São Pedro, no município de Vitória (ES). *Ciênc Saúde Coletiva* [Internet], Rio de Janeiro, 2010 July; 15(4): 2145-54. DOI: <http://dx.doi.org/10.1590/S1413-81232010000400028>

Publish in International Archives of Medicine

International Archives of Medicine is an open access journal publishing articles encompassing all aspects of medical science and clinical practice. IAM is considered a megajournal with independent sections on all areas of medicine. IAM is a really international journal with authors and board members from all around the world. The journal is widely indexed and classified Q1 in category Medicine.